

NASA/CR-1998-

206773

INTERIM  
70-76-012  
2017  
062-537

P3

**Interim Contractor Report #15**

1/01/98 - 1/31/98

**Physics of Boundaries and Their Interactions in Space Plasmas**

Dr. Nojan Omid  
Dr. Homayoun Karimabadi  
Dr. Dietmar Krauss-Varban

SciberNet, Inc.  
5414 Oberlin Drive, Suite 251  
San Diego, CA 92121

## **I. Large Scale Simulation of the Magnetopause:**

We have analyzed our run for southward IMF case in detail. We found a number of interesting features. The flow pattern at the magnetopause is quite complex, and is determined by the combined effect of the convection flow and the flows generated by the reconnection process. There are multiple reconnection sites and the thickness of the magnetopause varies significantly as one moves away from the subsolar point. In most instances, the reconnection site looks quite asymmetric and the fast flow can be generated almost entirely in one direction. Within each reconnection layer, the discontinuities that are formed are highly nonlocal in nature, and differ significantly from the usual discontinuities, such as rotational discontinuities that have been examined in isolation. We are currently writing up our findings for publication in the Geophysical Research Letters.

## **II. Three-dimensional Simulations**

We have analyzed a run with both the tearing mode and the Kelvin-Helmholtz instability. The evolution of the system is as follows. Initially, both KH and tearing mode start to grow. The growth for the tearing mode is almost all in the direction parallel to the background field, whereas the KH grows in the direction perpendicular to the field. This two modal property remains even in the highly nonlinear regime. What happens, however, is that tearing mode generates a core field, which is then amplified significantly by the modulations of the current layer caused by the KH instability. Further, both  $B_x$  and  $B_y$  grow in both the  $y$  and  $z$  directions. This is due to the fact that the tearing mode creates a current loop, which in turn becomes unstable to the KH in  $y$  as well as  $z$  direction. In the pure tearing case,  $B_x$  would only grow in the  $y$  direction. We are currently writing up the results for publication in the Geophysical Research Letters.

## **III. Inflow-Outflow Simulations:**

Using two-dimensional hybrid simulations of the magnetopause we have been able to show that: (1) nonlocal effects play a major role and affect the properties of the discontinuities associated with the reconnection layer. In particular, in contrast to previous studies, there are no 1-D equivalent of rotational discontinuities or intermediate shocks that are formed. In fact, if one tries to describe the discontinuities in terms of the usual discontinuities, one would be led to conclude that there are slow shocks in the magnetopause. But a more careful analysis, taking into account both the highly nonlocal and two-dimensional nature of the reconnection layer, reveals that the observed discontinuities are simply kinetic structures that have no simple fluid analogues. Further, we were able to demonstrate that the presence of the accelerated flows leads to the Kelvin-Helmholtz instability in the reconnection layer. Finally, in the case of multiple  $x$ -lines, we compared the cuts across the plasmoids with the observations of FTEs at the magnetopause. We showed how making different cuts through the plasmoids can lead to different field signatures. We are writing up our results for publication in the Geophysical Research Letters.

| REPORT DOCUMENTATION PAGE   |   |  | Form Approved<br>OMB No. 0704-0188                  |  |
|---|---|--|---|--|
| Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503  |   |  |   |  |
| 1. AGENCY USE ONLY (Leave Blank)  |   | 2. REPORT DATE<br>1/26/98                                  |   | 3. REPORT TYPE AND DATES COVERED<br>Interim Contractor Report, |
| 4. TITLE AND SUBTITLE<br>Physics of Boundaries and their Interactions in Space Plasmas  |   |  | 5. FUNDING NUMBERS<br><br>C: NAS5-96101             |  |
| 6. AUTHOR(S)<br>Dr. Nojan Omidi, Dr. Homayoun Karimabadi, and Dr. Dietmar Krauss-Varban   |   |  |   |  |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)<br>SciberNet, Inc.<br>5414 Oberlin Drive, Suite 251<br>San Diego, CA 92121   |   |  | 8. PERFORMING ORGANIZATION<br>REPORT NUMBER         |  |
| 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)<br><br>National Aeronautics and Space Administration<br>Goddard Space Flight Center<br>Greenbelt, MD 20771  |   |  | 10. SPONSORING / MONITORING<br>AGENCY REPORT NUMBER |  |
| 11. SUPPLEMENTARY NOTES   |   |  |   |  |
| 12a. DISTRIBUTION / AVAILABILITY STATEMENT<br>Unclassified-Unlimited  |   |  | 12b. DISTRIBUTION CODE                              |  |
| 13. ABSTRACT<br><br>This report describes the work done by SciberNet, Inc. during the month of January. During this time, we primarily worked on further analysis of the results presented at the AGU as well as writing them up for publication. Using large scale simulations, we showed that the magnetopause during the southward IMF case is quite irregular with varying thickness, and has a complex flow pattern owing to the nonlinear effects of the convective flow superimposed on the flows generated in the reconnection layer. We used inflow-outflow boundary conditions to examine the kinetic nature of the discontinuities that are formed in the reconnection layer and concluded that nonlocal effects play a major role in the formation of such discontinuities and can alter their properties from the usual structures expected from 1-D simulations or from fluid theories. Finally, we used our 3-D simulations to examine the nonlinear interaction of the tearing mode with the Kelvin-Helmholtz instability. We showed that this interaction leads to the generation of a large core field which is observed both in the magnetotail as well as the magnetopause. |   |  |   |  |
| 14. SUBJECT TERMS<br>Foreshock, bow shock, magnetotail, reconnection, kinetic simulations   |   |  | 15. NUMBER OF PAGES<br>1                            |  |
|   |   |  | 16. PRICE CODE                                      |  |
| 17. SECURITY CLASSIFICATION<br>OF REPORT<br>Unclassified  | 18. SECURITY CLASSIFICATION<br>OF THIS PAGE<br>Unclassified | 19. SECURITY CLASSIFICATION<br>OF ABSTRACT<br>Unclassified | 20. LIMITATION OF ABSTRACT<br><br>UL                |  |